



Armed Forces College of Medicine AFCM



Blood

White Blood Cells (Leukocytes)

Prof. Dr. Manal Hassan Moussa

Ass. Prof. Dr. Samaa Kamar

Intended Learning Objectives (ILO)



By the end of this lecture the student should
be able to:

- **Correlate** the structure of WBCs to its function
- **Interpret** the defective structure of the WBCs in different diseases

Lecture Plan



1. Part 1 (3 min) Introduction to white blood cells
2. Part 2 (40 min) white blood cells structure and function
3. Part 3 (3 min) Summary
4. Lecture Quiz (4 min)

Blood



- = A special type of connective tissue in which the matrix is fluid (plasma).
- Average volume in adult : 5 liters

Components of blood


Plasma

- 1- **Water** (92% of plasma)
- 2- **Proteins**
- 3- **Others:**
(Electrolytes, Nutrients, Respiratory gases, Waste)

Blood elements 45%

- 1- Red blood corpuscles (RBCs) = Erythrocytes
- 2- **White blood cells (WBCs) = Leucocytes**

Immunology and Blood Module




Leukocytes (White Blood Cells)



- **4500-11,000 /mm³ (μL)**
- **Colorless in fresh conditions / white if packed.**
- **True cells** having nuclei and organelles.
- **Spherical while suspended in blood plasma, but they become amoeboid and motile after**

to perform various activities related to immunity






Leukocytes (White Blood Cells)



Granulocytes

- **2 major types of granules (specific & non-specific granules)**
- **Life span of only a few days.**


Neutrophils **Eosinophils** Basophils




Agranulocytes

- **Lack specific granules but contain non-specific azurophilic granules.**
- **Life span: up years**

Lymphocytes



Monocytes




2 Major types of granules:

1. Specific granules:

Neutral
(neutrophils)

Acidic
(eosinophils)


Basic
(basophils)



2. Non-specific (Azurophilic) granules

- = **Lysosomes**
- Affinity for **methyl azure stain (blue-purple)**
- Present in **all leucocytes.**

Neutrophils



http://medcell.med.yale.edu/histology/blood_bone_marrow_lab/neutrophil.php

Neutrophils

Number: 60-70%

Size: 12-15 μm in diameter

Life span: short-lived cells (6-8hr)

L.M.: N: segmented deeply stained multi-lobed (2-5) interconnected by delicate chromatin strands.
(polymorph nuclear leucocytes =



“Band neutrophil”: immature neutrophil with horse shoe shape-like nucleus (< 1-2% in a blood film)



Barr body

**= *Drumstick-like appendage on one of the lobes
of the nucleus***

In 3% of neutrophils in peripheral blood films of

E.M.:

N: highly condensed chromatin

C: glycogen, small rER, Golgi, few mitochondria

2 Types of granules:

1- 1ry Azurophilic granules:


- larger (0.5 μm) / rounded / more electron dense
- = primary lysosomes
- Secrete myeloperoxidase, lysozyme

2- Specific granules:

- Smaller / less electron dense
- More numerous
- Secrete ECM degrading enzymes as collagenase, bactericidal


3- Tertiary granules :zymes.

- Contain metalloproteinase as gelatinase to help migration of neutrophils through the CT.



Function:

- o **Phagocytosis** of foreign organisms and destruction of bacteria
- o Neutrophils are inactive and spherical while circulating but **at areas of infection, neutrophils are first to arrive** (become amoeboid and leave the blood vessel by diapedesis) and become active and pursue the bacteria by chemotaxis.
- o Neutrophils **die** after **MARGINATION** **formation**.



Pus:
is accumulation of
dead leucocytes,
bacteria &
extracellular fluid)

Abnormal **Neutrophil** count

- **Neutrophilia (+++):**
Occur in bacterial infection

“Shift to the left”

++ percentage of band neutrophils

- **Neutropenia (- - -):**
Results in liability to bacterial infection


A 6 years old boy complained of dysphagia. His tonsils were red with white spots. Doctor asked for CBC. What do you expect?

Neutrophilia



https://www.researchgate.net/publication/315349776_Differential_diagnosis_of_tonsillitis_tonsillar_detritus_accumulation_and_tonsillar_keratin_cysts/figures?lo=1&utm_source=google&utm_medium=organic

Eosinophil




Eosinophil

Number 1-4% of total leucocytic count

Size: 12-15 μm in diameter

Life
8-10
days

L.M.: N: bilobed, less densely stained



than that of neutrophils

C: large retractile **acidophilic**
granules

E.M.:

N: highly condensed chromatin

C: small rER, Golgi, few mitochondria



2 Types of granules:

1- 1ry Azurophilic granules Non-specific lysosomes

2- Specific granules:

- large oval
- Have electron dense crystalloid core “internum” surrounded by a less electron dense matrix” externum”
- The crystalloid core = **major basic protein + peroxidase+**
(Major basic protein accounts for their acidophilic app
- The externum = large amount of arylsulfatase, histamin

prim



- Major basic protein, peroxidases and neurotoxins are cytotoxic to parasites



<https://gfycat.com/informalgoldenkudu>



Functions:

- 0 **Phagocytosis of antigen-antibody complexes**
- 0 **Secrete **histaminases** (inactivates histamine produced by mast cells).**
- 0 **Discharge their granules contents of on the surface of **parasitic worms** killing them**


Anti-allergic & Anti-parasitic

Abnormal **Eosinophil** count

- **Esinophilia (+++):**

Parasitic infection (Ex: bilharziasis)

Allergy (ex., asthma, urticaria and eczema,




- **Esinopenia (- - -): Corticosteroids**



https://www.zazzle.com/eosinophil+ts_hirts

Basophil



Basophil

Number:

<1% of total leucocytic count

Size:

10-12 μm in diameter



L.M.:

N: irregular lobed and usually obscured by granules

C: contains large dark basophilic granules

The granules are **metachromatic** (stained purple with toluidine blue)

Due to the presence of heparin and other sulfated GAGs.



E.M.:

N: ...

C: small rER, Golgi, few mitochondria


2 Types of granules:

1- 1ry Azurophilic granules:

- Primary lysosomes

2- Specific granules:

- large / more irregularly shaped
- Contain heparin, histamin, leukotrienes and eosinophil chemotactic factor,





Functions:

- Involved in **allergic** reaction...(receptors for Ig-E)
- Secrete their granular components in response to certain Ag/Ab reaction

Allergy

Comparison between Granulocytes



Basophils	Eosinophils	Neutrophils	
1%>	1-4%	60-70%	
µm 10-12	µm 12-15	µm 10-12	size
Irregular	Bilobed	Segmented multilobed ;2-5 dark lobes connected by delicate chromatin - <u>immature band neutrophil</u> . < 1-2%.	LM: nucleus
large dark basophilic gr. metachromatic	large retractile acidophilic granules	light neutral granules	Cytoplasm
small Golgi, few mitochondria -Azurophilic: lysosomes -Specific granules: heparin, histamine, eosinophilic chemotactic factor	small Golgi, few mitochondria -Azurophilic: lysosomes -Specific granules: electron dense crystalloid core internum major basic proteins , neurotoxin, peroxidase Histaminase	small Golgi , few mitochondria a-Azurophilic granules: lysosomes contain hydrolytic enz. , b-Specific granules: Small, numerous contain collagenase, c-Tertiary granules: gelatinase	EM



Agranulocytes


lack specific granules

But


contain azurophilic granules
(lysosomes)

Monocytes


Lymphocytes



Monocyte



Agranulocytes



Lymphocyte

Lymphocytes



Number **20-25%** of total leucocytic count.

Size: divided according to size into:


a-Small lymphocytes: about **7 μm** in diameter, **most numerous.**

c- Large lymphocytes

Functionally divided into:

- 1- **B**-lymphocytes (B-cells) 15%,
- 2- **T**-lymphocytes (T-cells) 80%
- 3- **Null** cells 5%

They have same morphology but can be differentiated by their **surface markers**



Small Lymphocytes




Size: 7 μm in diameter

L.M.:

N: - **large** occupying most of the cell,
- **Rounded**, shows a slight indentation.
- **Dense**.

C: a **thin rim** around the nucleus.

light basophilic with little azurophilic granules.



Larger lymphocytes have:

- Larger, slightly indented nuclei
- More cytoplasm that is slightly basophilic, with a few azurophilic gr.

Lymphocytes




E.M.:


Nucleus: rich in heterochromatin

The Cytoplasm contains:

- Abundant **ribosomes**.
- Few mitochondria, small Golgi complex, little rER.
- Few lysosomes (**azurophilic granules**)



d
Source: Mescher AL: Junqueira's Basic Histology: Text and Atlas, 12th Edition: http://www.accessmedicine.com
Copyright © The McGraw-Hill Companies, Inc. All rights reserved.






Functions:

- In blood stream, **No** function
- In connective tissue, they play a role in **immu**

B-lymphocytes T-lymphocytes

↓
Humoral

↓
Cytotoxic



https://sola.ai/i_t_r_on/t-lymphocyte-a-type-of-white-blood-cell-that-kills-cancer-1031553673


B-Lymphocyte function




Humoral-mediated immune response

When B lymphocytes recognize an antigen, they proliferate and differentiate

1- Plasma cells:
secrete antibodies



<http://apps.sanidadanimal.info/cursos/immunology-old/tercero2.htm>



2-Memory cells:

Re-exposure to same
antigen

**rapid and more
extensive**

production of antibodies;





T-Lymphocyte functions



Cellular-mediated immune response

1- Cytotoxic T-Lymphocytes:

they bind to the surface of foreign cells as viral infected cells & kill them




https://sola.ai/i_t_r_on/t-lymphocyte-a-type-of-white-blood-cell-that-kills-cancer-1031553

2-T-helper cells:

numerous factors → lymphocytes

activation of B-

and cytotoxic T-lymphocytes.



<http://leavingbio.net/human-defence-system/>

3-T-suppressor cells:

modulate extent of immune response and suppress production of antibodies

4- T memory cells: long lived,

responsible for a more rapid and extensive secondary immune response.

Null Lymphocyte function





- **Natural killer cells (NK)** that can kill foreign cells; virally infected, tumor cells
- Secrete **interferon γ** .

Lymphocyte Identification



**B and T lymphocytes are distinguished by detecting
“surface markers”
using immunohistochemical methods**



- **T-lymphocytes**: have T cell **receptors**,
T- cell receptors
CD molecules
- **B-lymphocytes**: have surface **immunoglobulins**.
- **Null cells**: have no surface markers for T nor B lymphocytes.




Compare between B and T Lymphocytes



	B lymphocytes	T lymphocytes
Percent	15%	80%
Origin	Bone marrow	Bone marrow
Site of maturation	Bone marrow	Thymus
Surface markers	Immunoglobulins	T-cell receptors Cluster of differentiation markers (CD2, CD3, CD4, CD8, and CD28)
Life span	Few months	Many years
Function	Humoral immune response	-Cell mediated immune response (T cytotoxic) -T-helper -T-suppressor



B-Cell Development




Monocytes



Number: 3-8% of total leucocytic count

Size: 12-20 μm . (largest blood cell).

L.M.:




Nucleus:

- **kidney-shaped**, deeply indented or a wide horse-shoe shaped with 2 nucleoli.
- **Eccentric** in position.
- Less condensed

Cytoplasm:

abundant, pale with **fine pinkish-purple azurophilic granules**.




Monocyte

Monocytes



E.M.:

- **Numerous lysosomes**
- rER, prominent **Golgi complex** near the indentation of the kidney-shaped nucleus
- Mitochondria.
- **Pinocytotic vesicles**
- Irregular processes



Monocyte Function




1- Phagocytosis:



Circulating monocytes can **phagocytose** bacteria and antigen antibody complexes in blood stream.

In the connective tissue they differentiate into macrophages.


2- Antigen presenting:



<https://gifimage.net/wp-content/uploads/2017/11/fagocytosis-gif-9.gif>



<https://gifimage.net/fagocitose-gif-4/>



<https://gifimage.net/fagocitose-gif-5/>

Lecture Quiz

Lecture Quiz



The cells that increase in suppurative tonsillitis are characterized by

- A-It is the largest leucocyte
- B-Granules with electron dense core
- C-Kidney shaped nucleus
- D-Azurophilic granules

In acute inflammation, first line of defense includes

- A-Eosinophils
- B-Basophils
- C-Neutrophils
- D-Lymphocytes

Lecture Quiz



A 45 year old patient presented with hematuria and was diagnosed as bilharziasis. His blood shows an increase in a leukocyte whose granules are characterized by:

- A-Secrete histamine
- B-Stain metachromatically
- C-Secrete collagenase
- D-Have electron dense internum

Which of the following is an antigen presenting cell?

- A- Basophils
- B- Neutrophils
- C- Monocytes
- D- Eosinophils

Lecture Quiz



Match the following cells with their contents

1- Neutrophils

a. Granules with electron dense core

2- Basophils

b. Granules containing collagenase

3- Eosinophils

c. Secrete histamine

4- Monocytes

d. Numerous lysosomes

SUGGESTED TEXTBOOKS



- 1. Junqueira`s Basic Histology; Text and Atlas. 14th edition 2016.**
- 2. Histology atlas and test: Michael H. Ross and Wojciech Pawlina, 7th edition, 2015.**



Thank
you

